

MCP

①

* Anxiety symp may occur in the following except:

- Hypoglyc
- Pheochromocytoma
- Hypoerthyroidism
- Conversion disorder
- AV prolapse

* Psychiatric exam includes:

- Mental state
- physical
- General
- Neuro
- All of above (✓)

* Delusions can be best defined as:-

- false fixed belief that meets specific (✓)
- perceptual misinterpretation

Representative

- a view point able to be changed when convincing evidence

* Depression occurs in except

- Parkinson's ds
- panic ϕ
- Hypothyroidism
- MV prolapse (\checkmark)

NB All GIT ϕ may present $\hat{=}$ depression

* All types of delusions except:

- Persecut
- Grandiosity
- Infidelity
- Thought echo (\checkmark)

* Mania is ~~ecc~~ by except

- Euphoria
- Nihilistic delusions clear! ϕ (\checkmark)
- Hyperactivity
- Delusions of grandiosity

* Schizophrenia is ccc by all except

- Delusions
- Visual hallucinations
- impaired insight
- ~ memory (✓)
- (-ve) symptoms

* Best tx of ADHD is

- Amphetamine (✓) → "stim the Cx that inhibits lower centers"
- MAOIs
- ECT
- Benzodiazepines

* Antidep drugs include all except

- MAOIs
- SSRI
- Serpasil (✓) → "it's ant HTN"
- TCAs

* Extra pyramidal side effects of antipsychos.

- Rigidity & tremors (✓)
- Euphoria
- obsessions
- Dementia

* Risk factors of delirium include:

- Dehydrat
- ICU
- delirium (✓)
- None

NB Delirium mostly in both extremes of age
 ↳ Start nocturnal

* Tendency to ↑ the dose of psychoactive substance to get the same effects is of

- Habituation
- saturation
- Tolerance (✓)
- abuse
- synergism

* The following substance \rightarrow dependence

Except:

- Tryptizole (✓)
- Cocaine
- Amphetamine
- Benzodiazepines
- Caffeine

* Social phobia is ccc by:

- Fear of animals
- ~ heights
- ~ talking in public (✓)
- ~ enclosed spaces

* Pharmacological ttt of Bipolar:

- Carbamazepine
- No valproate
- Lithium
- All (✓)

* ECT is at of

- Anxiety
- Depression (✓)

* Causes of excitement except

- Mania
- Dilemma
- Social phobia (✓)
- Paranoid schizo
- Catatonia (✓) *Q's up!*
(high over-kill!)

* BZD & anticonvulsants exert
action through:

- Muscarinic
- Dopaminergic
- GABAergic (✓)

* Diff bet obsessions & compulsions:

- Acts, which the pte has to do to
alleviate their anxiety (✓)
- Ideas

* ECT used in ALL except

Panic disorders (✓)

* Pharmacotherapy of depi:

- BZD

- SSRIs (✓)

- BB

* " of schizophrenia

- Phenothiazine (✓)

- Amphetamines

- BZD

* Suicide is more common in, except,

- post partum psychosis

- severe depression

- previous suicidal attempt

- conversion disorder (✓)

* Antipsychotic act:

- Blocking dopaminergic (✓)

Page _____
Date _____

⑤

* Depression is CCC by except,

- ↓ self confidence (✓)
- Guilt feeling
- lack of interest
- suicidal

* Acute dystonia dt antipsychotic
can be tx by:

- Anti hist
- Anti chol
- BZD
- All (✓) "it's an emergency"
- None

* Post traumatic stress disorder, except

- The stressor is major & out of normal human experience

Flash back

- No marked occup. impairment (✓)
- Avoidance symptoms

* The following can produce psychiatric symp.

- CS
- BB
- INF
- OCP
- ALL (✓)

* Symp of GAD :

- Apprehension
- tremors
- Tachy
- most of time
- All (✓)

* Schizo is CCC by all except

- Delusions
- And Hallucination
- Disorientate to T, P, P (✓)
- formed thought disorder

Problem solving

① IB\$ → want of \$
→ Or shopping أريد المال

Ans ⇒ Hypochondriasis

② cat of The Blue → fear, sense of Death
⇒ Panic disorder

③ Urinary retention by
(xatropine)
⇒ TCA

④ M, 55 yr, experience easy fatigue
lack of interest in everyday activities
difficult to sleep
⇒ Thyro tox.

5. Bizarre behaviour \Rightarrow schizophrenia

6. ♀ clonic conv., clothes torn

No cyanosis, No tongue biting

No urinary incontinence

\Rightarrow Hysterical fits

7. aggression, disorientation to T, BP,

\Rightarrow Dementia

تذكروا
"الأذن، العين، القلب، الأذن، العين، القلب"
الجنون